

# GetValidTest

## Try Before You Buy

Download a free sample of any of our exam questions and answers

- ✓ 24/7 customer support, Secure shopping site
- ✓ Free One year updates to match real exam scenarios
- ✓ If you failed your exam after buying our products we will refund the full amount back to you.

Select a vendor... ▾

Select an exam... ▾

Your email address

 Free Download



<http://www.getvalidtest.com>

GetValidTest - Valid test study guide, get certification

**Exam : BAS-011**

**Title : IBM SPSS Statistics Level 1  
v2**

**Vendors : IBM**

**Version : DEMO**

NO.1 With IBM SPSS Statistics you can assign one of three levels of measurement to a variable. Which

is the correct list of all three levels?

- A. Nominal - Ordinal - Interval
- B. Categorical - Interval - Ratio
- C. Nominal - Ordinal - Scale
- D. Nominal - Interval – Scale

Answer: C

NO.2 If you request an aggregated dataset or data file in the Aggregate procedure, the number of cases

in the new aggregated file is equal to what?

- A. Number of cases in the original data file
- B. Number of aggregated summary variables
- C. Number of categories of the variables specified in the Break Variables list
- D. Number of cases that you specified in the Aggregate Data dialog box

Answer: C

NO.3 In the Aggregate data procedure<sup>1</sup> if you specify more than one Break Variable, \_\_\_\_\_ in the

Break Variables list will determine case order.

- A. Variable size
- B. Variable label
- C. Variable name
- D. Variable order

Answer: D

NO.4 If you use the IBM SPSS Statistics Select Cases dialog to conduct analysis on a subset of cases and you want to keep all cases in the same dataset, which option is appropriate?

- A. Copy unselected cases to a new dataset
- B. Copy selected cases to a new dataset
- C. Delete unselected cases
- D. Filter out unselected cases

Answer: D

NO.5 What type of graph would you use to see the distribution of a scale variable?

- A. Bar chart
- B. Pie chart
- C. Histogram
- D. Scatter plot

Answer: C

NO.6 In the output of the CROSSTABS procedure, large residuals indicate agreement between the data

and the null hypothesis of independence.

- A. True
- B. False

Answer: B

NO.7 When you read data stored in an Excel file via File ... Open ... Data, you may need to specify which options? (Choose three.)

- A. Read variable names from the first row of data
- B. Read a Range of cells
- C. Specify the Worksheet to read/import
- D. Number of Worksheets to read/import

Answer: A,B,C